Ghiath Tayeb, M.D. Board Certified Gastroenterology Reem Jarbou, D.O. Board Certified Gastroenterology Seina Farshadsefat, D.O

Phone: 248-651-0800 Fax: 248-651-7341

Outpatient Instructions for Colonoscopy

Plenvu Preparation

Patient's Name		
Hospital/Facility		
Date of Procedure		
Time to Arrive		

* Arrival time is subject to change. The facility will call you the day before the procedure with the exact arrival time. *

One Day Prior to the Procedure:

- Begin clear liquid diet. (See diet instructions on the following page)
- Try to drink lots of liquids to help with the cleansing and to prevent dehydration. You may continue the clear liquid until eight hours prior to the procedure.

Day of Procedure:

- Eight hours prior to the procedure do not eat or drink anything, including water.
- No medications may be taken in the morning before the procedure except heart and blood pressure medication (take with a *small* sip of water). Any other medications should be taken after the procedure is over.
- You may NOT work or drive the day of the procedure. You must have a driver that will remain in the facility during the procedure and drive you home or the test will be cancelled.
- If patient is taking weight loss/diabetic injection must stop injections as follows: if taking daily injection, hold injection the day of procedure. If taking weekly injections hold injection one week prior to procedure.

You may continue taking your regular medication during the preparation, with the following exceptions:

- STOP taking Aspirin, Motrin, and Arthritis medication 7 days prior to the procedure. If needed, you may take Tylenol products.
- **STOP** taking Iron (check multi-vitamin ingredients for iron), Coumadin, Warfarin, or any blood thinners (Plavix, Ticlid, Effient, or Pradxa) 3 days prior to the procedure.

If you have any questions or concerns, please call the office at (248)-651-0800.

If you are unable to keep your appointment, please give at least <u>two business days'</u> notice. A \$100 fee will be assessed for cancellations made after that time. Thank you.

Patient Instructions for Plenvu

- 1. Pick up preparation kit at your pharmacy.
- 2. Begin the clear liquid diet the day before your procedure.
- 3. The evening before you procedure at 5 pm:
 - A) Pour Dose 1 packet in clear container.
 - B) Add water to top of container and mix.
 - C) Drink all the liquid in the container within 30 minutes.
 - D) Fill the container with any clear liquid of your choice and drink.
- 4. The evening before you procedure at **8 pm**:
 - E) Pour Dose 2 (POUCH A <u>AND</u> POUCH B TOGETHER) into clear container.
 - F) Add water to the top of container and mix.
 - G) Drink all the liquid in the container.
 - H) Fill the container with any clear liquid of your choice and drink.

^{*}There can only be a 2 hour gap between the two doses.

^{*}Each dose must be drunk within a half an hour of being mixed with water.

^{*}You can refrigerate the solution.

CLEAR LIQUID DIET INSTRUCTIONS

Day to	begin vour	clear liquid diet	•
	7 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	1-1	

Start your clear liquid diet when you wake up in the morning. No food, milk, or milk product (including powdered creamers) for the entire day and until the procedure has been completed.

No red, green, or blue liquids due to the dye colors! Yellow, orange, or clear liquids are permitted.

Foods included on the clear liquid diet:

Beverages: Carbonated beverages (soda & diet soda), coffee, tea, Gatorade,

apple juice, orange Hi-C, white grape juice or white cranberry

juice.

Desserts: Jello, popsicles, frozen ices/slushes, Italian ices, freeze pops.

Soups: Bouillon, broth (chicken or beef).

Sweets: Honey, sugar, plain hard sugar candy (yellow or orange only).

Condiments: Salt, pepper.

Fruits: Strained (not just pulp free) pineapple, orange, or grapefruit

juice.